

ANTIBACTERIAL ACTIVITY OF *Xetospongia* sp. AND *Spongia* sp. SPONGE EXTRACT AGAINST PATHOGENIC BACTERIA FROM THE WATERS OF PAHAWANG ISLAND, LAMPUNG

Aktivitas Antibakteri Ektrak Spons Jenis *Xetospongia* sp. dan *Spongia* sp. terhadap Bakteri Patogen dari Perairan Pulau Pahawang, Lampung

Anma Hari Kusuma*, Berta Putri, Eko Efendi, Tiara Safitri, Siska Hidayah

Marine Science Study Programe, Universitas of Lampung, Lampung

Prof. road. Dr. Ir. Sumantri Brojonegoro No.1, Gedong Meneng, Rajabasa, Bandar Lampung

*Corresponding author: anma.hari@fp.unila.ac.id

(Received October 29th 2025; Accepted December 28th 2025)

ABSTRACT

Sponges are marine benthic invertebrates from the Porifera phylum that have bioactive compounds that can function as antibacterials. The purpose of this study was to compare the extracts of sponges of the *Xetospongia* sp. and *Spongia* sp. against pathogenic bacteria. This study was conducted in May-November 2025. Samples of sponges of the *Xetospongia* sp. and *Spongia* sp. taken from the waters of Pahawang Island were extracted using methanol, ethyl acetate and hexane. The results showed that the morphological characteristics the *Xetospongia* sp. had a rough body surface with a rounded shape and reddish color while the *Spongia* sp. type had a soft body surface with crusts and a dark black color. The yield of sponge extract using ethyl acetate was greater than using methanol and hexane. The antibacterial activity of sponge extracts against pathogenic bacteria of the *Xetospongia* sp. used hexane while the *Spongia* sp. used methanol.

Keywords: Sponge, Yield, Antibacterial Activity

ABSTRAK

Spons adalah hewan avertebrata bentik laut dari Filum Porifera yang memiliki senyawa bioaktif yang dapat berfungsi sebagai antibakteri. Tujuan penelitian ini adalah membandingkan antara ekstrak spons jenis *Xetospongia* sp. dengan *Spongia* sp. terhadap bakteri patogen. Penelitian ini dilakukan pada bulan Mei-November 2025. Sampel spons jenis *Xetospongia* sp. dan *Spongia* sp. diambil dari perairan Pulau Pahawang diekstrak menggunakan metanol, etil asetat dan heksana. Hasil penelitian menunjukkan ciri morfologi memiliki permukaan tubuh kasar dengan bentuk membulat dan warna kemerahan sedangkan jenis *Spongia* sp. memiliki permukaan tubuh lembut dengan mengerak dan warna hitam gelap. Rendemen ekstrak spons menggunakan etil asetat lebih besar dibandingkan dengan menggunakan metanol dan hexana. Aktivitas antibakteri ekstrak spons terhadap bakteri patogen spons *Xetospongia* sp. menggunakan hexana sedangkan spons *Spongia* sp. menggunakan metanol.

Kata Kunci: Spons, Rendemen, Aktivitas Antibakteri

INTRODUCTION

Infectious diseases are a significant health problem. They are highly dynamic and can spread directly or indirectly from one person to another. The mortality rate from these diseases remains relatively high (Ristori *et al.*, 2024). Every year, infectious diseases can cause significant mortality. Treatment for infectious diseases typically involves antibiotics. Inappropriate and uncontrolled antibiotic use accelerates the emergence of antibiotic-resistant bacterial strains, commonly referred to as Multi-Drug Resistance (MDR) (Mancusso *et al.*, 2021). This has become a complex health issue. Concerns about the management of infectious diseases caused by MDR strains have prompted numerous studies exploring new antibiotic compounds. Exploration of bioactive compounds from the seabed has been conducted in various regions and has yielded numerous compounds with potential for use as chemotherapeutic agents (Karthikeyan *et al.*, 2022). Indonesia's vast waters offer promising marine resources for the exploration of various new chemical compounds

Sponges are marine benthic invertebrates from the phylum Porifera. Their habit of living in extreme conditions has led to their unique defense system, called secondary metabolites (Maslin *et al.*, 2021). These secondary metabolites are used for self-defense and adaptation mechanisms to extreme environments, competition for space, defense against predation, inhibition of attachment organisms, and defense against other organisms (Lee *et al.*, 2021). The results of this metabolism are bioactive compounds. These bioactive compounds are compounds from the terpenoid, alkaloid, polypeptide, polyeptide, and polyphenol groups (Perdicaris *et al.*, 2013). These bioactive compounds can function as antibacterials, antifungals, antivirals, antitumor, anticancer, antifouling agents, and enzyme inhibitors (Rajendran, 2019). One sponge that possesses bioactive compounds is a sponge from the Demospongiae class, namely the species *Xetospongia* sp. and *Spongia* sp. The aim of this study was to compare the effectiveness of sponge extracts from *Xetospongia* sp. and *Euspongia* sp. against pathogenic bacteria.

METHODS

Time and Location Research

This research was conducted from May to November 2025 in the waters of Pahawang Island, Pesawaran Regency, Lampung Province (Figure 1). Laboratory analysis was conducted at the Oceanography Laboratory, Department of Fisheries and Marine Sciences, Faculty of Agriculture, University of Lampung.




Figure 1. Map of research location

Tool and Materials

The tools used in this study include SCUBA (Self-Contained Underwater Breathing Apparatus) diving equipment, masks, gloves, scissors, 600 ml bottles, cool boxes, rulers, knives, Erlenmeyer flasks, rotary evaporators, analytical scales, separating funnels, measuring cups, petri dishes, autoclaves, tweezers, spirit burners, magnetic stirrers, dropper pipettes, micro tubes, hot plates, vortexes, spatulas, laminar air flow, test tubes, underwater camera incubators, and GC-MS while the materials used are sponge identification books, sponge samples, stain pathogenic bacteria *Escherichia coli*, distilled water, methanol, ethyl acetate, hexane, chloramphenicol, disc paper, filter paper and tissue.

Research Procedures

Sponge Sampling

Sampling was conducted using SCUBA diving equipment. Sponge samples were collected and cut using a knife, amounting to 1 kg of each type. They were then wrapped in plastic, placed in a cooler, and transported to the laboratory.

Sponge Ectraction

Extraction was carried out using the principle of maceration in methanol, ethyl acetate, and hexane solvents. 50 g sample was placed in a 100 ml solution of methanol, ethyl acetate, and hexane solvents and left for 48 hours, then filtered using filter paper and a vacuum pump (Kusuma *et al.*, 2023). The filtrate obtained was then evaporated at 60 °C to obtain a crude extract in the form of a paste. The crude sponge extract was dried at room temperature until dry, weighed, labeled, and stored at 4 °C.

Bacterial Culture

Culture of the pathogenic *E. coli* bacteria was aseptically inoculated into a test tube containing 5 ml of Nutrient Broth (NB) medium. The medium was incubated for 24 hours at 37°C. Then, the medium was inoculated again from the NB medium into Nutrient Agar (NA) medium, incubated for another 24 hours, and stored as a stock of the bacteria.

Antibacterial Test

Bacterial effectiveness testing was conducted using the Kirby-Bauer or paper disk method, which involves observing and measuring the inhibition zone of sponge extract. A 1 ml inoculum of pathogenic *E. coli* bacteria was aseptically taken from NB medium and poured into a Petri dish containing NA medium. Then, a paper disk approximately 0.5 mm in diameter was dipped into sponge extract at specific concentrations with each solvent. The disc was placed using tweezers on top of the NA medium containing pathogenic *E. coli* bacteria and incubated at 37°C for 24 hours. All media were then incubated for another 24 hours at 37°C, and the diameter of the inhibition zone was observed and measured until the third day (Kusuma *et al.*, 2023).

Data Analysis

The yield data was analyzed by comparing the weight of the obtained extract with the weight of the dry raw materials used in the extraction process. Antibacterial activity data was analyzed by calculating the diameter of the inhibition zone, expressed as the clear area around the paper discs for each sponge type and solvent type.

RESULT

Sponge Identification

Xetospongia sp. has a globular, reddish-colored barrel-shaped body with a rough surface. *Xetospongia* sp. is a type of sponge with a body shape resembling a large vase or barrel, commonly called a barrel sponge. The color of this sponge varies depending on the species and environment, but is typically pink, reddish-brown, purple, or grayish-brown. The outer surface of this sponge is rough or uneven in texture, sometimes with small protrusions. Ostia are found throughout the body surface. This sponge has a large osculum at the top, through which water exits after being filtered. The diameter of the osculum can be quite large and conspicuous. The body skeleton is made of spongin, a protein, and spicules are made of silica. (Hasan *et al.*, 2022; Khodzori *et al.*, 2023). *Spongia* sp. has a dark, encrusting body with a smooth surface. *Spongia* sp. is a type of sponge with an irregular shape that can crust, branch, or resemble mounds, so this sponge can be said to lack clear body symmetry. This sponge's color varies, generally light yellow, brown, or yellowish gray depending on the aquatic environment. Ostia of this sponge are scattered throughout the body. The body skeleton is spongin from protein, but the spicules are without silica. (Prastiyanto *et al.*, 2022).




Figure 2. Marine sponge (a) *Xetospongia* sp. and (b) *Spongia* sp. at research location

Sponge Extract Yield

Extraction results showed varying values in methanol, ethyl acetate, and hexane solvents. Sponge extract yields at the research site can be seen in Table 1.

Table 1. Sponge extract yield

Marine Sponge	Morphology	Yield (%)		
		Methanol	Ethyl Acetate	Hexane
<i>Xetospongia</i> sp.	Globular	2,24	8,33	2,61
<i>Spongia</i> sp.	Encrusting	9,79	10,08	1,81

The yield of *Xetospongia* sp. sponge extract using methanol was 2,24%, ethyl acetate was 8,33%, and hexane was 2,61%, while *Spongia* sp. sponge using methanol was 9,79%, ethyl acetate was 10,08%, and hexane was 1,81%. The yield of *Xetospongia* sp. sponge extract using methanol was 1,01-5,53%, ethyl acetate was 4,43-8,56%, and hexane was 3,34-9,61% (Swantara *et al.*, 2019; Ibrahim *et al.*, 2021; Azemi *et al.*, 2019) while *Spongia* sp. sponge using methanol was 1,01-5,53%, ethyl acetate was 4,43-8,56%, and hexane was 3,34-9,61% (Swantara *et al.*, 2019; Ibrahim *et al.*, 2021; Azemi *et al.*, 2019). using methanol of 2,34-7,78%, ethyl acetate of 5,67-8,89%, hexane of 1,81-9,87%. (Krishnan & Keerthi, 2016; Yang & Liang 2024; Tai *et al.*, 2022). The yield of sponge extract using ethyl acetate is greater than using methanol and hexane. The yield of sponge extracted using ethyl acetate is higher at 10,08% compared to hexane at 1,81% (Chen *et al.*, 2022; Majali *et al.*, 2015).

Antibacterial Activity of Sponge Extract Against Pathogenic Bacteria

The antibacterial activity of a bioactive compound is depicted by the clear zone of inhibition formed around the paper disc. The antibacterial activity of sponge extract against pathogenic bacteria at the study site is shown in Table 2.

Table 2. Antibacterial activity of sponge extract against pathogenic *E. coli*

Marine Sponge	Morphology	Inhibition Zone (mm)			
		Methanol	Ethyl Acetate	Hexane	Chloramphenicol (Control)
<i>Xetospongia</i> sp.	Globular	0,6	0	17,3	18,4
<i>Spongia</i> sp.	Encrusting	5,2	4,2	3,8	20,2

The activity of sponge extract against pathogenic bacteria *E. coli* obtained inhibition zone for *Xetospongia* sp. using methanol of 0,6 mm and hexane of 17,3 mm while *Spongia* sp. using methanol of 5,2 mm, ethyl acetate of 4,2 mm and hexane of 3,8 mm. The activity of sponge extract against pathogenic bacteria *E. coli* obtained inhibition zone for *Xetospongia* sp. using methanol of 3 mm and hexane of 25 mm (Edny et al, 2020) while the antibacterial activity of *Spongia* sp. sponge against pathogenic bacteria *E. coli* with methanol of 8 mm and hexane of 3 mm (Kamaruding et al, 2020). The antibacterial activity of *Xetospongia* sp. sponge extract is found in hexane solvent while *Spongia* sp. sponge in methanol solvent. This is also the same as what was conveyed in (Hartiadi et al., 2020; Kalinggi et al, 2017) where the antibacterial activity of the sponge type *Xetospongia* sp. used hexane while the sponge *Spongia* sp. used methanol (Yang & Liang 2024).

DISCUSSION

The solubility of bioactive substances is highly dependent on the solvent's ability to dissolve a substance. The level of polarity is a very important factor in determining the value of a yield because solvents with different polarities will dissolve different bioactive substances. The yield of sponge extracts from each type of solvent shows varying results. Methanol is an alcohol compound of a carbon group bonded to three hydrogen atoms and one hydroxyl group. Hexane is an alkane compound with a straight chain with six carbon atoms. Ethyl acetate is an ester compound of acetic acid and ethanol. The yield of sponge extracts using ethyl acetate solvent is greater than that of methanol and hexane for each type of sponge as a whole because ethyl acetate has properties between hydrophobic and hydrophilic. Ethyl acetate has a medium polarity that allows for the production of yield extracts. Various forms of bioactive substances in the form of yield extracts in an organism have complex structures that are neither completely polar nor non-polar but are in the middle spectrum so that ethyl acetate solvent is more effective in producing yields compared to methanol and hexane.

The antibacterial activity of sponge extract against the pathogenic bacteria *E. coli* for the sponge type *Xetospongia* sp. using hexane solvent was greater than that of methanol and ethyl acetate. Conversely, for the sponge type *Spongia* sp. antibacterial activity using methanol solvent was greater than that of ethyl acetate and hexane. This proves that the sponge type *Xetospongia* sp. has bioactive compounds as non-polar antibacterials, while the sponge type *Spongia* sp. has bioactive compounds as polar antibacterials. The bioactive compounds of the sponge type *Xetospongia* sp. Non-polar compounds include terpenoid groups such as xestolone, xestosaprol, xestobergsterol, steroid groups such as xesterol derivatives, 24-methylcholesta-5,22-dien-3 β -ol, xesterol, fatty acid groups such as palmitic acid, stearic acid and oleic acid, and carotenoids (Ke et al., 2024 & Gamal et al., 2016). Polar bioactive

compounds of *Spongia* sp. sponges include alkaloid groups such as spongidine a–e, saponin groups such as spongoside, peptide groups such as spongipeptin A–D, amino acid groups such as spongin acid and phenol groups such as spongiaphenol (Maximo *et al.*, 2016).

CONCLUSION

Sponges found in the waters of Pahawang Island for the type *Xestospongia* sp. have a rounded (globular) reddish body with a rough body surface while the type *Spongia* sp. has a dark encrusting body with a soft body surface. The yield of sponge extract using ethyl acetate is greater than using methanol and hexane. The highest antibacterial activity of *Xestospongia* sp. sponge extract is found in hexane solvent compared to methanol and ethyl acetate while *Spongia* sp. sponge in methanol solvent compared to ethyl acetate and hexane. *Xestospongia* sp. sponge extract with hexane solvent and *Spongia* sp. with methanol solvent can be used as candidates for antibacterial against pathogenic bacteria *E. coli*.

ACKNOWLEDGEMENT

We would like to express his gratitude to fellow lecturers and students at the University of Lampung who assisted in the research and writing of this scientific paper. He also thanks the Institute for Research and Community Service (LPPM) of the University of Lampung (Unila) for funding this research through the Unila BLU DIPA basic research scheme

REFERENCE

- Azemi, N. A., Ismail, N., Baka, L. A., & Muhammad, T. S. T. (2019). Sterol Composition and Anti-Atherosclerosis Effects of *Xestospongia muta* Extract by Increasing Transcriptional Activity of SR-B1 Promoter. *International Journal of Pharmaceutical Sciences and Research*, 10(6), 2892–2897. <https://doi.org/10.13040/IJPSR.0975-8232.10.6.2892-97>
- Chen, W. C., Wang, S. W., Li, C. W., Lin, H. R., Yang, S. C., Chu, Y. C., Lee, T. H., & Chen, J. J. (2022). Comparison of Various Solvent Extracts and Major Bioactive Components from *Portulaca oleracea* for Antioxidant, Anti-Tyrosinase, and Anti-Glucosidase Activities. *Antioxidants*, 11(2), 398-415. <https://doi.org/10.3390/antiox11020398>
- Edny, G., Rebecca, J., Sheryl, C., Crystalia, A. A., & Hartiadi, L. Y. (2022). A Review on the Antimicrobial Properties of Giant Barrel Sponge *Xestospongia* sp. *Indonesian Journal of Life Sciences*, 2(2), 96–112. <https://doi.org/10.54250/ijls.v2i2.55>
- Gamal, A. A. E., Mansoor, S., Shaala, L. A., Alhadaid, A. M., Al-Said, M. S., Ashour, A. E., Kumar, A., Abdel-Kader, M. S., Abdel-Mageed, W. M., & Youssef, D. T. A. (2016). Cytotoxic Compounds from the Saudi Red Sea Sponge *Xestospongia testudinaria*. *Marine Drugs*, 14(5), 82-91. <https://doi.org/10.3390/md14050082>
- Hartiadi, L. Y., Edny, G., Rebecca, J., Sheryl, C., & Crystalia, A. A. (2020). A Review on the Antimicrobial Properties of Giant Barrel Sponge *Xestospongia* sp. *Indonesian Journal of Life Sciences*, 2(2), 1–10. <https://doi.org/10.54250/ijls.v2i2.55>
- Hasan, T., Sofiana, M. S. J., Apriansyah, Hartanti, L., Lestari, D., Safitri, I., & Nurdiyansyah, S. I. (2022). Antioxidant Activity of Crude Protein of *Xestospongia* sp. Sponge from Spermonde Sulawesi. *Jurnal Biologi Tropis*, 22(4), 1291–1296. <https://doi.org/10.29303/jbt.v22i4.4402>
- Ibrahim, A., Rudiyanisya, & Alimuddin, A. H. (2021). Karakterisasi Struktur Lipid dari Spons *Xestospongia testudinaria*. *Indonesian Journal of Pure and Applied Chemistry*, 4(1), 30–39. <https://doi.org/10.26418/indonesian.v4i1.45537>
- Kalalinggi, S. Y., Soekamto, N. H., Firdaus, Appa, F. E., & Irmawan, M. (2023). Phytochemical Screening n-Hexane Extract of Sponge *Xestospongia testudinaria* from

- Spermonde Archipelago. *Journal of Biotropical Research and Nature Technology*, 1(2), 25–45. <https://doi.org/10.52850/borneo.v1i2.8992>
- Kamaruding, N. A., Ismail, N., & Sokry, N. M. (2020). Identification of Antibacterial Activity with Bioactive Compounds from Selected Marine Sponges. *Pharmacognosy Journal*, 12(3), 493–502. <https://doi.org/10.5530/pj.2020.12.76>
- Ke, M. L., Zhang, Z. R., Li, S. W., Zeng, Y. B., Su, M. Z., & Guo, Y. W. (2024). Bioactive Steroids with Structural Diversity from the South China Sea Soft Coral *Lobophytum* sp. and Sponge *Xestospongia* sp. *Marine Drugs*, 22(1), 36-48. <https://doi.org/10.3390/md22010036>
- Khodzori, F. A., Mazlan, N. B., Chong, W. S., Palaniyeloo, K., & Shah, D. M. (2023). Metabolites and Bioactivity of the Marine *Xestospongia* Sponges (Porifera, Demospongiae, Haplosclerida) of Southeast Asian Waters. *Biomolecules*, 13(3), 484-500. <https://doi.org/10.3390/biom13030484>
- Krishnan, K. A., & Keerthi, T. R. (2016). Analyses of Methanol Extracts of Two Marine Sponges, *Spongia officinalis* var. *ceylonensis* and *Sigmadocia carnosa* from Southwest Coast of India for Their Bioactivities. *International Journal of Current Microbiology and Applied Sciences*, 5(2), 722–734. <https://doi.org/10.20546/ijcmas.2016.502.081>
- Kusuma, A. H., Kurniawan, K., & Gipari, N. (2023). Aktivitas Antibakteri Ekstrak Karang Lunak terhadap Bakteri Patogen *Escherichia coli*. *Jurnal Perikanan*, 13(3), 925–934. <https://doi.org/10.29303/jp.v13i3.614>
- Kusuma, A. H., Nugraha, A. H., & Shasika, A. B. (2023). Antibacterial Activity of Seagrass Extract against Pathogen Bacteria of *Escherichia coli* Strain Multi Drug Resistance (MDR). *Jurnal Biologi Tropis*, 23(4), 171–178. <https://doi.org/10.29303/jbt.v23i4.5486>
- Lakshmi, V., & Shukla, P. K. (2014). Antimicrobial Activity of Marine Sponge *Spongia officinalis* var. *ceylonensis* Dendy. *Journal of Marine Biology & Oceanography*, 3(3), 1–4. <https://doi.org/10.4172/2324-8661.1000131>
- Majali, I., Qaralleh, H., Idid, S. Z., Saad, S., Susanti, D., & Althunibat, O. Y. (2015). Potential Antimicrobial Activity of Marine Sponge *Neopetrosia exigua*. *Journal of Basic and Applied Research*, 1(1), 1–13. <https://doi.org/10.48550/arXiv.1912.05323>
- Maximo, P., Ferreira, L. M., Branco, P., Lima, P., & Lourenco, A. (2016). The Role of *Spongia* sp. in the Discovery of Marine Lead Compounds. *Marine Drugs*, 14(8), 139. <https://doi.org/10.3390/md14080139>
- Prastiyanto, M. E., Kartika, A. I., Darmawati, S., & Radjasa, O. K. (2022). Bioprospecting of Bacterial Symbionts of Sponge *Spongia officinalis* from Savu Sea, Indonesia for Antibacterial Potential against Multidrug Resistant Bacteria. *Biodiversitas Journal of Biological Diversity*, 23(2), 1118–1124. <https://doi.org/10.13057/biodiv/d230256>
- Saroinsong, M. S., Kandou, F. E. F., Papu, A., & Singkoh, M. F. O. (2014). Uji Daya Hambat Ekstrak Metanol Beberapa Jenis Porifera terhadap Bakteri *Escherichia coli* dan *Staphylococcus aureus*. *Jurnal MIPA*, 3(2), 129–133. <https://doi.org/10.35799/jm.3.2.2014.5989>
- Swantara, I. M. D., Rita, W. S., Suartha, I. N., & Agustina, K. K. (2019). Anticancer Activities of Toxic Isolate of *Xestospongia testudinaria* Sponge. *Veterinary World*, 12(9), 1434–1440. <https://doi.org/10.14202/vetworld.2019.1434-1440>
- Tai, C. J., Ahmed, A. F., Chao, C. H., Chang, F. R., Huang, Y. M., Yen, C. H., Hwang, T. L., & Sheu, J. H. (2022). Spongenolactones A–C: Bioactive Pentacyclic Diterpenes from a Red Sea *Spongia* sp. *Marine Drugs*, 20(8), 498-510. <https://doi.org/10.3390/md20080498>
- Yang, Q. B., & Liang, L. F. (2024). *Spongia* Sponges: Unabated Sources of Novel Secondary Metabolites. *Marine Drugs*, 22(5), 213-224. <https://doi.org/10.3390/md22050213>